



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Mr. Maynard has reported (*Birds of E. N. A.*, 1879, p. 168) seeing a Canada Jay in Newtonville "in early summer" about 1875, but Mr. Mann's bird seems to be the first that has been actually taken in Massachusetts.—WILLIAM BREWSTER, *Cambridge, Mass.*

**A Second Nest and Eggs of *Picicorvus columbianus* taken in Colorado.**—Mr. Denis Gale, of Gold Hill, Boulder Co., Colorado, writes me that he found a second nest of Clark's Nutcracker, containing three fresh eggs, on April 16, 1889. He found the nesting site first on March 12, noticing one of the birds trying to break off a small twig from a dead tree, and watching the direction it took afterwards. The bird which came from behind him flew high over his head, and after waiting for some time he noticed both birds flying to and from a certain point fully five or six times. After a careful search he finally found a few twigs lying upon a horizontal limb in the dense top of a small scrubby pine-tree about twelve feet high and six inches in diameter. On visiting the place again a week later, at his approach one of the birds flew off a neighboring tree uttering his warning note, but upon inspection he found no changes or augmentation of the twigs, and concluded therefrom that a new nesting site had been selected, and when he visited the spot a week later still with the same results his conclusion seemed to be verified and he made an exhaustive search within a radius of a mile of this point, but all to no purpose, seeing neither nests nor birds. Four weeks later, on April 16, passing close to the place where the birds had commenced building first, he stopped to look once more at the site first selected by this pair of birds, and much to his surprise discovered a bulky nest in place of the few twigs first noticed, with the female on it and covering three eggs. The nest was placed about nine feet from the ground, and resembles the one taken in 1888 in every particular; perhaps it is a little more bulky still. The eggs bear a close resemblance to the first set found by Mr. Gale, excepting that the markings are possibly a little more decided and numerous. They measure  $1.34 \times .90$ ,  $1.37 \times .91$ , and  $1.39 \times .92$  inch. Mr. Gale's first nest, containing three fresh eggs also, was found on March 5, 1888, six weeks earlier and probably an unusually early case.—CHARLES E. BENDIRE, *Washington, D. C.*

**Bullock's Oriole in Maine.**—Mr. Manly Hardy writes me that a male Bullock's Oriole (*Icterus bullocki*) was shot "a few miles from Bangor, Maine, about the middle of November, 1889, and sent in the flesh to Mr. Crosby, the well-known Bangor taxidermist, by whom it was mounted." Mr. Hardy has lately examined the bird and compared it with a Western specimen, from which it differs only in being "a little more of a canary color."

This capture adds a species to the New England list as well as to the fauna of Maine. It also affords still another example of the curious fact that most of the Western and Southern birds which occur in New England as rare or purely accidental stragglers, are found in late autumn or early winter.—WILLIAM BREWSTER, *Cambridge, Mass.*